20

CLAIMS

We claim:

1. A method for determining the differences between the molecular interactions of two different mixtures of molecules, comprising:

labeling a first mixture of molecules and a target mixture of molecules; introducing said first mixture of molecules to a combinatorial library of solid phase supports;

incubating said combinatorial library with said first mixture of molecules; performing a first marking step to mark those of said solid phase supports that have a molecule of said first mixture bound to them;

introducing said target mixture of molecules to said combinatorial library; incubating said combinatorial library with said target mixture of molecules; obtaining a first image showing as marked those of said solid phase supports

that have a molecule of said first mixture bound to them;

performing a second marking step to mark those of said solid phase supports

that have a molecule of said target mixture bound to them;

obtaining a second image showing as marked those of said solid phase supports that have a molecule of said target mixture bound to them;

and

20

a molecule of said target mixture bound to them, wherein said third image is created by comparing said first image and said second image.

2. The method of Claim 1, further comprising:

isolating one of said solid phase supports identified in said third image; and determining the chemical structure of a ligand on one of said isolated solid phase supports.

- 3. The method of Claim 1, wherein said first mixture of molecules is a protein extract from normal cells and said target mixture of molecules is a protein extract from cancer cells.
- 4. The method of Claim 1, wherein said combinatorial library is a one-bead-one-compound peptide library.
- 5. The method of Claim 1, wherein said labeling is performed by biotinylation.
- 6. The method of Claim 5, further comprising:

before said first marking step is performed, incubating said combinatorial library with a solution of streptavidin-alkaline phosphatase conjugate; and,

after said incubating said combinatorial library with said target mixture of molecules is performed, but before said obtaining said first image is

20

performed, incubating said combinatorial library with a solution of streptavidin-alkaline phosphatase conjugate.

- 7. The method of Claim 1, wherein said labeling is performed by the use of an antigen and corresponding antibody.
- 8. The method of Claim 1, wherein said first and said second marking steps are performed by incubating said combinatorial library in a solution of 5-bromo-4-chloro-3-indolyl-phospate.
- 9. The method of Claim 1, wherein said combinatorial library is immobilized in a support matrix before said first image is obtained.
- 10. The method of Claim 1, wherein said first and said second images are graphical images, and said third image is created by comparing said first and said second images on a pixel-by-pixel basis.
- 11. The method of Claim 10, wherein said first image is image "A," said second image is image "B," and said third image is created by applying the formula (B-A)/A on a pixel-by-pixel basis.
- 12. A method for determining the differences between the molecular interactions of two different mixtures of molecules and identifying a ligand specific for a molecule in one of the mixtures, comprising:

labeling a first mixture of molecules and a target mixture of molecules;

introducing said first mixture of molecules to a combinatorial library of solid phase supports;

incubating said combinatorial library with said first mixture of molecules; performing a first marking step to mark those of said solid phase supports

introducing said target mixture of molecules to said combinatorial library;

that have a molecule of said first mixture bound to them;

obtaining a first image showing as marked those of said solid phase supports

incubating said combinatorial library with said target mixture of molecules;

that have a molecule of said first mixture bound to them;

performing a second marking step to mark those of said solid phase supports

that have a molecule of said target mixture bound to them;

obtaining a second image showing as marked those of said solid phase

supports that have a molecule of said target mixture bound to them;

creating a third image identifying those of said solid phase supports that have

a molecule of said target mixture bound to them, wherein said

third image is created by comparing said first image and said

second image;

isolating one of said solid phase supports identified in said third image; and determining the chemical structure of a ligand on one of said isolated solid

phase supports.

5

10

- 13. The method of Claim 12, wherein said first and said second images are graphical images, and said third image is created by comparing said first and said second images on a pixel-by-pixel basis.
 - 14. The method of Claim 13, wherein said first image is image "A," said second image is image "B," and said third image is created by applying the formula (B-A)/A on a pixel-by-pixel basis.
 - 15. A method for identifying a ligand specific for a molecule in one of two different mixtures of molecules, comprising:

labeling a first mixture of molecules and a target mixture of molecules; introducing said first mixture of molecules to a combinatorial library of solid phase supports;

incubating said combinatorial library with said first mixture of molecules; performing a first marking step to mark those of said solid phase supports that have a molecule of said first mixture bound to them;

introducing said target mixture of molecules to said combinatorial library; incubating said combinatorial library with said target mixture of molecules; obtaining a first image showing as marked those of said solid phase supports

performing a second marking step to mark those of said solid phase supports that have a molecule of said target mixture bound to them;

that have a molecule of said first mixture bound to them;

8141/9886/EAN/588344.WPD;

-38-

File No. 8141/9886

ø

obtaining a second image showing as marked those of said solid phase supports that have a molecule of said target mixture bound to them; creating a third image identifying those of said solid phase supports that have a molecule of said target mixture bound to them, wherein said third image is created by comparing said first image and said second image;

isolating one of said solid phase supports identified in said third image; and determining the chemical structure of a ligand on one of said isolated solid phase supports.

16. A method for identifying a ligand specific for a target molecule, comprising: labeling a target molecule;

incubating a combinatorial library of solid phase supports with a label binder;

performing a first marking step to mark those of said solid phase supports that have a molecule of said label binder bound to them; introducing said target molecule to said combinatorial library; incubating said combinatorial library with said target molecule; obtaining a first image showing as marked those of said solid phase supports that were marked in said first marking step;

20

20

performing a second marking step to mark those of said solid phase supports that have a target molecule bound to them;

obtaining a second image showing as marked those of said solid phase supports that have a target molecule bound to them;

creating a third image identifying those of said solid phase supports that have a target molecule bound to them, wherein said third image is created by comparing said first image and said second image;

isolating one of said solid phase supports identified in said third image; and determining the chemical structure of a ligand on one of said isolated solid phase supports.

- 17. The method of Claim 16, wherein said labeling is performed by biotinylation, and further, wherein said label binder is streptavidin-alkaline phosphatase conjugate.
- 18. The method of Claim 16, wherein said labeling is performed by the use of an antigen and corresponding antibody.
- 19. A method of screening a combinatorial library, comprising:

labeling a target molecule;

incubating a combinatorial library of solid phase supports with a label binder;

performing a first marking step to mark those of said solid phase supports that have a molecule of said label binder bound to them;

introducing said target molecule to said combinatorial library;

incubating said combinatorial library with said target molecule;

obtaining a first image showing as marked those of said solid phase supports that were marked in said first marking step;

performing a second marking step to mark those of said solid phase supports that have a target molecule bound to them;

obtaining a second image showing as marked those of said solid phase supports that have a target molecule bound to them; and creating a third image identifying those of said solid phase supports that have a target molecule bound to them, wherein said third image is created by comparing said first image and said second image.

20. The method of Claim 19, further comprising:

isolating one of said solid phase supports identified in said third image; and determining the chemical structure of a ligand on one of said isolated solid phase supports.